



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Laplace assumed such law of increase of density from surface to center as would make the differential equation for the ellipticities of the strata

$$\frac{d^2}{da^2} \left[\varphi(a)e \right] = \frac{6}{a^2} \varphi(a)e + 3a^2 e \frac{d\rho}{da} \quad (1)$$

integrable.

By assuming that the last term of the second member of equation (1) is a function of a , Pratt, in his "Figure of the Earth," deduced the result

$$\rho = Q \frac{\sin qa}{a},$$

in which ρ = the density, Q and q are constants, and a = radius of shell considered as spherical.

The constants q and Q are found from the conditions that the density at the surface is 2.75, and the mean density of the Earth is twice that of the surface.

By means of the densities thus found, Pratt deduces the masses of the nucleus, and four concentric shells, the radius of the nucleus and the thickness of each shell being equal to one fifth of the Earth's radius.

The masses in order from the surface are, 0.3372, 0.3257, 0.2237, 0.0981, 0.0153, the mass of the Earth being unity.

From these values the masses of the parts whose radii are R , $\frac{4}{5}R$, $\frac{3}{5}R$, $\frac{2}{5}R$ and $\frac{1}{5}R$ are found to be 1.0000, 0.6628, 0.3371, 0.1134, 0.0153.

Taking the formula for attraction at the surface, each mass considered a sphere, $A = m \div R^2$, we get in order from the surface, $\frac{1}{25}(1.0000)$, $\frac{1}{16}(0.6628)$, $\frac{1}{9}(0.3371)$, $\frac{1}{4}(0.1134)$, 0.0153, or, 0.0400, 0.0414, 0.03746, 0.0284, 0.0153.

If this is the true law, the force of attraction increases from the surface till we reach a point perhaps one-third distant from surface to center, from that point the force diminishes and becomes zero at the center.

SOLUTION OF THE PROBLEM PROPOSED BY MR. HARVILL AT THE CLOSE OF HIS SOLUTION ON PAGE 56.

BY THE EDITOR.

LET A represent the center of the sphere, radius r , BCD a tri-rectangular spherical triangle, and $AEFGHPCI$, the solid removed from this portion of the sphere in making the first hole.

Draw BPh , DPi and CP , arcs of great circles respectively through BP , DP and CP , and draw HPj , IPk , small circles thro' P with poles at B , D .

The solid $A E F G H P I C$ consists of a square pyramid, $A E F G P$ plus two similar and equal triangular spherical pyramids, $C B P A$ and $C D P A$, minus two equal sectors $B P I E$ and $D P H G$, of two equal spherical segments whose height is $B E = D G$.

Putting $2x$ for the width of the chisel, we have $A E = E F = x$, and consequently $F P$, the altitude of the square pyramid $= \sqrt{(r^2 - 2x^2)}$; hence the solidity of the pyramid $A E F G P = \frac{1}{3}x^2\sqrt{(r^2 - 2x^2)}$.

The base of the spherical pyramid $B C P A$, is the area of the spherical triangle $B C P = S$.

To find S , we have $\angle C = 45^\circ$, $\angle B = \sin^{-1}[x \div \sqrt{(r^2 - x^2)}]$, whence we find $\angle P = \sin^{-1}[\sqrt{(r^2 - 2x^2)} \div \sqrt{(2r^2 - 2x^2)}] + 90^\circ$; therefore the sum of the angles of the spherical triangle $B C P$ is

$$\sin^{-1}\left[\frac{r+2x}{(r+x)\sqrt{2}}\right] + 135^\circ;$$

$$\therefore S = \left\{ \sin^{-1}\left[\frac{r+2x}{(r+x)\sqrt{2}}\right] - 45^\circ \right\} \frac{S}{90^\circ},$$

where S' is the area of the spherical trian. $A B C = \frac{1}{8}$ the surface of the sphere $= \frac{1}{2}\pi r^2$. Hence the solidity of the two equal spherical pyramids is

$$\frac{\frac{1}{3}\pi r^3}{90^\circ} \left\{ \sin^{-1}\left[\frac{r+2x}{(r+x)\sqrt{2}}\right] - 45^\circ \right\}.$$

The solidity of the sectors to be subtr. is $\frac{\frac{1}{8}(4r^3 - 6r^2x + 2x^3)}{360^\circ} \cdot \sin^{-1} \frac{x}{\sqrt{(r^2 - x^2)}}$; hence $\frac{1}{8}$ the solidity removed in cutting the first hole through the sphere is

$$\frac{x^2\sqrt{(r^2 - 2x^2)}}{3} + \frac{\frac{1}{3}\pi r^3}{90^\circ} \left\{ \sin^{-1}\left[\frac{r+2x}{(r+x)\sqrt{2}}\right] - 45^\circ \right\} - \frac{\frac{1}{8}(4r^3 - 6r^2x + 2x^3)}{360^\circ} \cdot \sin^{-1} \frac{x}{\sqrt{(r^2 - x^2)}},$$

and consequently the part of the sphere removed in making the two holes will be 16 times the above $-8x^3$, the part common to both holes. Therefore

$$\frac{16x^2\sqrt{(r^2 - 2x^2)}}{3} + \frac{\frac{1}{3}\pi r^3}{90^\circ} \left\{ \sin^{-1}\left[\frac{r+2x}{(r+x)\sqrt{2}}\right] - 45^\circ \right\} - 8x^3 - \frac{\frac{1}{8}(4r^3 - 6r^2x + 2x^3)}{360^\circ} \cdot \sin^{-1} \frac{x}{\sqrt{(r^2 - x^2)}} = \frac{2\pi r^3}{3}.$$

By solving this transcendental equation the value of x will be found.

